

Notable invertebrates associated with wet woodland

Snails and slugs (Mollusca)		
<i>Perforatella rubiginosa</i>	RDB3	
<i>Phenacolimax major</i>	Na	
<i>Acicula fusca</i>	Nb	
<i>Leiostyla anglica</i>	Regionally local	
<i>Spermodea lamellata</i>	Regionally local	
Spiders and allies (Arachnida: Araneae and Pseudoscorpiones)		
<i>Carorita paludosa</i>	RDB2	a small money-spider
<i>Centromerus levitarsis</i>	RDB2	a money-spider
<i>Agraecina striata</i>	Nb	a hunting spider
<i>Diplocephalus protuberans</i>	Nb	a money-spider
<i>Gongylidiellum murcidum</i>	Nb	a money-spider
<i>Hygrolycosa rubrofasciata</i>	Na	a wolf-spider
<i>Hypomma fulvum</i>	Na	a money-spider
<i>Maro sublestus</i>	Nb	a money-spider
<i>Maso gallicus</i>	Na	a small money-spider
<i>Notioscopus sarcinatus</i>	Nb	a money-spider
<i>Porrhomma oblitum</i>	Nb	a money-spider
<i>Saloca diceros</i>	Nb	
Centipedes (Chilopoda)		
<i>Lithobius muticus</i>	Nb	
Woodlice (Isopoda)		
<i>Trichoniscoides albidus</i>	Nb	
Stoneflies (Plecoptera)		
<i>Nemoura dubitans</i>	N	
Dragonflies and damselflies (Odonata)		
<i>Somatochlora metallica</i>	N	Brilliant Emerald dragonfly
<i>Gomphus vulgatissimus</i>	N	Club-tailed dragonfly
<i>Cordulia aenea</i>		Local Downy Emerald dragonfly
True bugs (Hemiptera)		
Heteroptera		
<i>Cymus aurescens</i>	Nb	
<i>Globiceps flavomaculatus</i>	Nb	
<i>Orthotylus virens</i>	Nb	
<i>Pachycoleus waltli</i>	Nb	
Leafhoppers, planthoppers, froghoppers, treehoppers & cicadas (Auchenorrhyncha)		
<i>Aphrophora alpina</i>	Nb	a large brown leafhopper
Flies (Diptera)		
Snail-killing flies, picture-wing flies, grass flies and allies (Acalyptrata)		
<i>Eurygnathomyia bicolor</i> (Pallopteridae)	RDB1	
<i>Homoneura interstincta</i> (Lauxaniidae)	RDB3	
<i>Tanypeza longimana</i> (Tanypezidae)	RDB2	
<i>Elachiptera austriaca</i> (Chloropidae)	N	
<i>Pelidnoptera nigripennis</i> (Phaeomyiidae)	N	Larva parasitoid of millipedes

Typhamyza bifasciata (Anthomyzidae) N

Soldier flies, bee flies, robber flies & allies (Larger Brachycera)

<i>Stratiomys chamaeleon</i>	RDB1	Clubbed General
<i>Oxycera analis</i>	RDB2	Dark-winged Soldierfly
<i>Oxycera pardalina</i>	N	Hill Soldierfly
<i>Oxycera pygmaea</i>	N	Pygmy Soldierfly
<i>Ptiolina obscura</i>	N	Black-fringed Moss-snipefly
<i>Stratiomys potamida</i>	N	Banded General
<i>Spania nigra</i>	N	Liverwort Snipefly

Hoverflies (Syrphidae)

<i>Parasyrphus nigritarsis</i>	BAP Priority	RDB1
more or less characteristic of wet woodland		
<i>Chalcosyrphus eunotus</i>	RDB2	
<i>Cheilosia nebulosa</i>	RDB3	
<i>Cheilosia uviformis</i>	[Status not known]	
records seem to indicated Priority Habitat Wet woodland, but very few records		
<i>Meligramma guttatum</i>	N	
<i>Orthonevra geniculata</i>	N	
<i>Sphegina verecunda</i>	N	
<i>Xylota abiens</i>	N	
<i>Xylota tarda</i>	N	

Craneflies (Tipulidae)

<i>Lipsothrix nervosa</i>	BAP Priority	
<i>Lipsothrix errans</i>	BAP Priority	
<i>Lipsothrix nigristigma</i>	BAP Priority	
<i>Ellipteroides alboscutellata</i>	RDB1	
<i>Erioptera nigripalpis</i>	RDB3	
<i>Dicranomyia nigristigma</i>	RDBK	
<i>Scleroprocta pentagonalis</i>	RDB3	
<i>Dicraniomyia aquosa</i>	N	
<i>Dicraniomyia stigmatica</i>	N	
<i>Molophilus corniger</i>	N	
confined to calcareous seepages in woodland		
<i>Molophilus variispinus</i>	N	
largely confined to calcareous seepages in upland woodland		
<i>Scleroprocta sororcula</i>	N	
wet woodlands only		
<i>Paradelphomyia fuscula</i>	N	
confined to wet woodland		
<i>Paradelphomyia nielseni</i>	N	
confined to wet woodland		
<i>Thaumastoptera calceata</i>	N	
wet woodland only		
<i>Tipula yerburyi</i>	N	

Blowflies, dung flies, flesh flies and allies (Calyprata)

<i>Helina pulchella</i> (Muscidae)	RDB3	
<i>Hemyda vittata</i> (Tachinidae)	RDB3	
<i>Piezura boletorum</i> (Fanniidae)	RDBK	
<i>Subclytia rotundiventris</i> (Tachinidae)	RDB3	
<i>Elocera delecta</i> (Tachinidae)	N	
<i>Fannia corvina</i> (Fanniidae)	N	

<i>Fannia gotlandica</i> (Fanniidae)	N
<i>Fannia speciosa</i> (Fanniidae)	N
<i>Lophosia fasciata</i> (Tachinidae)	N
<i>Phaonia siebecki</i> (Muscidae)	N
<i>Phaonia zugmayeriae</i> (Muscidae)	N
Beetles (Coleoptera)	
Leaf beetles (Chrysomelidae) (Leaf beetles)	
<i>Cryptocephalus decemmaculatus</i>	BAP Priority RDB 2
<i>Agelastica alni</i>	RDB K
<i>Luperus flavipes</i>	Nb
<i>Lythraria salicariae</i>	Nb
Weevils (Curculionoidea)	
<i>Melanapion minimum</i>	BAP Priority
<i>Orchestes testaceus</i>	BAP Priority
<i>Dorytomus majalis</i>	RDB IK
<i>Is nigratarsis</i>	Na
<i>Cossonus linearis</i>	Na
<i>Cossonus parallelepipedus</i>	Nb
<i>Cryptorhynchus lapathi</i>	Nb
<i>Magdalis carbonaria</i>	Nb
<i>Scolytus ratzeburgi</i>	Nb
Rove beetles and allies (Staphylinidae/Scydmaenidae/Silphidae)	
<i>Oxypoda nigrocincta</i>	RDBI
limited to floodplain woodland in the vicinity of slow-flowing lowland rivers	
<i>Scaphisoma boleti</i>	Nb
<i>Omalium rugatum</i>	N
<i>Oxytelus fulvipes</i>	Na
largely confined to carr and swampy areas associated with riverine fen	
<i>Gabrius velox</i>	Nb
<i>Sepedophilus bipunctatus</i>	Nb
<i>Sepedophilus pedicularius</i>	N
<i>Sepedophilus testaceus</i>	N
<i>Gyrophaena joyi</i>	N
typical of wet woodland, especially that dominated by grey willow, but occasionally found in drier woodland types and unwooded fens	
<i>Gyrophaena lucidula</i>	N
<i>Bibloplectus pusillus</i>	N
<i>Carpelimus impressus</i>	Local
<i>Lathrobium elongatum</i>	Local
<i>Lathrobium impressum</i>	Local
<i>Bryaxis puncticollis</i>	Local
Long-horned beetles (Cerambycidae)	
<i>Obera oculata</i>	BAP Priority RDB1
<i>Lamia textor</i>	RDB1
<i>Gracilia minuta</i>	RDB2
<i>Aromia moschata</i>	Nb
<i>Saperda carcharias</i>	Na
<i>Phymatodes alni</i>	Nb
Click beetles (Elateroidea)	
<i>Diacanthous undulatus</i>	Nb

Caddis flies (Trichoptera)

<i>Ironoquia dubia</i>	RDB2
<i>Adicella filicornis</i>	RDB3
<i>Tinodes pallidulus</i>	RDB1
<i>Emodes articularis</i>	N
<i>Hydropsyche fulvipes</i>	N
<i>Phacopteryx brevipennis</i>	N
<i>Plectrocnemia brevis</i>	N
<i>Tinodes rostocki</i>	N
<i>Tinodes unicolor</i>	N
<i>Hydatophylax infumatus</i>	Local
<i>Lasiocephala basalis</i>	Local and regionally notable
<i>Limnephilus ignavus</i>	Local & regionally notable
<i>Trichostegia minor</i>	Regionally notable
<i>Lype reducta</i>	Common

Ants, bees and wasps (Hymenoptera: Aculeata)

<i>Pemphredon morio</i>	RDB3, Nb, Scarce
-------------------------	------------------